MultiDip

NEW





- Offers a tick-detaching effect.
- Controls nuisance, biting and face flies.
- Protects against screw worm infestation.



1,5 L 1000 L WATER

REPLENISH

3 to 1000 L

WATER







CYMIAZOLE

175 g/l

CYPERMETHRIN

25 g/l



1,5 L 1000 L WATER

MultiDip

Boost the spray wash by adding 200 ml of MultiDip after every 100 head of cattle dipped.

MultiDip

NOTE: It is not good spray race practice to carry dip wash over from one spraying to the next. Only make up sufficient wash for each spraying (± 3 litres per head plus the amount for the pump system).



15 ml 10 L WATER

Thoroughly wet the animal by spraying from below upwards so that the hair is lifted by the spray. Concentrate on the ears, under the tail and in the tail brush.

DIPPING INTERVALS:

TICKS, FLIES AND SCREW WORMS: Dip or spray weekly LICE: Dip or spray when necessary

Virbac RSA (Pty) Ltd (Reg. No. 1990/003743/07), Private Bag X115, Halfway House 1685, South Africa Tel: +27 12 657-6000 Fax: +27 86 226 8432 | za.virbac.com

MultiDip - Reg. No: G4444 (Act/Wet 36/1947); Botswana Reg. No: W130942; Zimbabwe Reg. No: 2019/80.16.12/9795 Composition: Cymiazole 175 g/l and Cypermethrin 25 g/l

Virbac