



Flukazole C

FOR ANIMAL USE ONLY

Flukazole C

Reg. No. G3533 (Act 36/1947)



A drench for the control of liver fluke, roundworms, lungworms and tapeworms in cattle and sheep

CAUTION

COMPOSITION:

Triclabendazole 12 % m/v
Oxfendazole 4,53 % m/v

STORAGE INSTRUCTION:

Store below 30 °C in a cool, dry place

WARNINGS:

- Do not slaughter animals for human consumption within 28 days of dosing.
- Do not dose lactating animals whose milk or milk products are intended for human consumption.
- Do not use less than 21 days before calving/lambing, in animals where milk or milk products from treated animals may be used for human consumption.
- Where accidental treatment with Flukazole C takes place within the 21 day period, milk must not be used for human consumption, or be supplied for processing for at least 21 days following treatment. Young animals fed this milk within 21 days of slaughter must not be used for human consumption.
- Containers and any residual contents should be disposed of safely in accordance with national waste disposal regulations. Empty containers should not be re-used for any other purpose.
- Dangerous to fish. DO NOT contaminate dams, rivers or streams with the drench or used container.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS:

- Wash hands after use.
- Avoid contact with eyes and skin. May cause irritation.
- Rinse the medicated drench guns/apparatus three times after use. Do not dispose of undiluted chemicals.
- Wash equipment thoroughly after use.

FIRST AID:

If swallowed, and if more than 15 minutes from a hospital, induce vomiting.

DIRECTIONS FOR USE: Use only as directed

Shake before use.

Cattle and Sheep

Dosage: 1 ml / 10 kg body mass, orally. This dose delivers 12 mg Triclabendazole per kg body mass and 4,53 mg Oxfendazole per kg body mass. Cattle in excess of 650 kg body mass should be dosed at 5 ml per 50 kg body mass. Dose the group according to the body mass of the heaviest animal in the group.

EFFICACY:

Roundworms

Flukazole C controls benzimidazole-sensitive mature and immature roundworms and lungworms.

Roundworms - Cattle	Immatures	Adult
Wireworm (<i>Haemonchus placei</i>)	*	*
Brown stomach worm (<i>Ostertagia ostertagi</i>)	*	*
Cattle bankrupt worm (<i>Cooperia</i> spp.)	*	*
Hookworm (<i>Bunostomum phlebotomum</i>)	*	*
Nodular worm (<i>Desophagostomum radiatum</i>)	*	*
Lungworm (<i>Dictyocaulus viviparus</i>)	*	*

Definition: * Control (≥ 90 % effective)

Roundworms - Sheep

	Immatures	Adult
Wireworm (<i>Haemonchus</i> spp)	*	*
Brown stomach worm (<i>Teladorsagia</i> spp)	*	*
Large mouth bowel worm (<i>Chabertia</i> spp)	*	*
Long-necked bankrupt worm (<i>Nematodirus</i> spp)	*	*
Lung worm (<i>Dictyocaulus</i> spp)	*	*
Bankrupt worm (<i>Trichostrongylus</i> spp)	*	*
Hook worm (<i>Gaigeria</i> spp)	*	*
White bankrupt worm (<i>Strongyloides</i> spp)	*	*

Definition: * Control (≥ 90 % effective)

Ovicidal: Roundworm eggs present in the animal at the time of dosing are prevented from hatching.

Cattle and sheep should not be placed on clean pasture until 6 hours after drenching.

Tapeworm: Flukazole C also controls tapeworm in cattle and sheep.

Milk tapeworm (*Moniezia* spp). Class 1 (100 % effective in 80 % of the treated animals).

Liver Fluke: Flukazole C is used for the treatment of susceptible early immature, immature and mature Liver Fluke.

Flukazole C controls the following


	Early immatures		Late immatures			Adult
	2 weeks	3 weeks	4 weeks	6 weeks	8 weeks	
Liver Fluke (<i>Fasciola hepatica</i>)	*		*	*	*	*
Giant Liver Fluke (<i>Fasciola gigantica</i>)		*		*		*

Definition: * Control (≥ 90 % effective)

The list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information contact your veterinarian.

REGISTRATION HOLDER

VIRBAC RSA (Pty) Ltd
(Reg No. 1990/003743/07)
Private Bag X115
Halfway House
1685
South Africa
Tel: +27 12 657 6000

Namibia Reg. No. V09/18.1.2/77 
Botswana Reg. No. W130659