

FOR ANIMAL USE ONLY

Phenix[®] CTC 20%

Reg. No. G2133 (Act 36/1947)

Pigs: As an aid in the control of diseases caused by chlortetracycline-sensitive organisms, e.g. atrophic rhinitis, enzootic pneumonia, *Actinobacillus (Haemophilus) pleuropneumonia*, necrotic enteritis.

Fowl: For the treatment and control of susceptible gastrointestinal and respiratory infections.

Cattle: Is recommended for oral administration via feed as an aid in the prevention and control of disease in beef cattle, caused by organisms susceptible to chlortetracycline.

Calves: An aid in the control of bacterial enteritis, primary or secondary bacterial pneumonia and other bacterial infections, particularly if the calf is subject to stress following housing changes, transportation or chilling.

■ STORAGE INSTRUCTIONS

Store below 25 °C in a dry place, protected from light.

■ COMPOSITION

Chlortetracycline 200 g (20 %) per kg.

■ WARNINGS

- Keep out of reach of children, uninformed persons and animals.
- Calves, cattle and pigs should not be slaughtered for human consumption during treatment or for 7 days thereafter.
- Fowls should not be slaughtered for human consumption during treatment or for 4 days thereafter.
- Do not administer to laying fowls.
- Do not use in breeding animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

■ PRECAUTIONS:

- Adequate measures must be taken not to create dust when incorporation of the product into feed is occurring. Persons handling Phenix CTC 20 % should do so in a mechanically ventilated area.
- Use dust mask and wear protective gloves and approved safety glasses during addition of Phenix CTC 20 % to feed.
- Wash exposed skin after handling and before eating.
- Do not smoke or eat during use.

■ DIRECTIONS FOR USE**Use only as directed**

Dosage and administration: For oral administration after incorporation in complete feed. To ensure thorough dispersal, Phenix CTC 20 % should be first mixed with 25 kg feed ingredients before incorporation in the final mix. Included in feed at a rate of 1,5 - 2 kg / ton of complete feed to supply the equivalent of 300 - 400 ppm dietary chlortetracycline.

Swine: Phenix CTC 20 % should be included in the feed to provide a dose of 20 mg chlortetracycline per kg body mass daily. This can generally be achieved by adding 2 kg Phenix CTC 20 % per ton of complete feed. This may be increased up to 6 kg per ton of complete feed at a veterinary surgeon's discretion to ensure adequate dosing of inappetent animals.

Fowls: The recommended dose is 20 - 60 mg chlortetracycline per kg body mass daily. This can generally be achieved by adding 1,5 - 2 kg Phenix CTC 20 % per ton of complete feed. This constitutes an average dose, but dosage levels may be varied between 100 ppm and 500 ppm according to the nature of and severity of infection, feed intake in relation to body mass and the appetite of diseased animals.

Treatment is recommended for 5 - 7 days for pigs, fowl, cattle and calves. Allow treated pigs, fowl, cattle and calves access only to medicated feed during treatment.

■ PRESENTATION

A brown granular meal for incorporation in animal feed in packs of 25 kg.

■ REGISTRATION HOLDER

VIRBAC RSA (Pty) Ltd
(Reg. No. 1990/003743/07)
Private Bag X115
Halfway House
1685
South Africa
Tel: +27 12 657 6000

Namibia Reg. No. V06/17.1.2/610 NSO