

FOR ANIMAL USE ONLY

Phenix® Oxytetracycline 20 % FG PMX

Reg. No. G2748 (Act 36/1947)

For the treatment of secondary bacterial infections following respiratory diseases in poultry and for pigs. Bacterial scours and pasteurella in calves and lambs. Gastroenteritis in pigs caused by bacteria sensitive to oxytetracycline and for the use against other bacterial infections as prescribed by a veterinarian.

■ STORAGE INSTRUCTIONS

Store at room temperature below 25 °C.

■ COMPOSITION

Oxytetracycline 20 % m/m.

■ WARNINGS

- **Withdrawal period:** Meat 7 days.
- Do not use in breeding or dairy animals.
- Do not use in laying hens.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

■ PRECAUTIONS:

- Not to be used after expiry date indicated on the bag
- Medicated feed must be used within three months after mixing.

■ DIRECTIONS FOR USE

Use only as directed

Mix the required mass of Phenix Oxytetracycline 20 % FG PMX thoroughly in a small quantity of complete feed (20 - 30 kg) before it is incorporated into the bulk of the total feed to be prepared.

Species	Phenix Oxytetracycline 20 % FG PMX (kg / 1 000 kg feed)	Oxytetracycline (g / 1 000 kg feed)	Treatment period (days)
Calves and lambs	0,25 - 1,5	50 - 300	7 - 14
Pigs	0,5 - 2	100 - 400	7 - 14
Poultry (broilers and breeders)	0,5 - 1	100 - 200	7 - 14

■ PRESENTATION

1 kg, 5 kg, 10 kg and 25 kg.

■ REGISTRATION HOLDER

VIRBAC RSA (Pty) Ltd
(Reg. No. 1990/003743/07)
Private Bag X115
Halfway House
1685
South Africa
Tel: +27 12 657 6000