

FOR ANIMAL USE ONLY

Promote®

Reg. No. G1247 (Act 36/1947)

TO IMPROVE THE RATE OF MASS GAIN AND FEED EFFICIENCY IN PIGS, LAYER BIRDS AND BROILERS.

■ **STORAGE INSTRUCTIONS**

Store in a cool, dry place.

■ **COMPOSITION**

Tylosin 10 % m/m. More than 80 % of tylosin is Factor A

■ **WARNINGS**

- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

■ **PRECAUTIONS**

- Not to be used after the expiry date indicated on the bag.
- Medicated feed must be used within 3 months after mixing.

■ **DIRECTIONS FOR USE****Use only as directed**

Mix appropriate mass of PROMOTE per metric ton of complete feed to obtain the recommended level. It is recommended that PROMOTE should be thoroughly mixed in a small quantity of feed (20 - 30 kg) before it is incorporated into the bulk of the total feed to be prepared.

Pigs:

| Pig mass (feed category) | PROMOTE (g/1 000 kg feed) | Tylosin (mg / kg feed) |
|--------------------------|---------------------------|------------------------|
| 11 - 18 kg (Creep) | 400 | 40 |
| 19 - 45 kg (Grower) | 200 - 400 | 20 - 40 |
| 46 kg (Finisher) | 200 | 20 |

Layer birds: Mix 220 g (22 mg / kg) to 550 g (55 mg / kg) PROMOTE per metric ton of feed, depending on the age of the flock and the *Mycoplasma gallisepticum* status of the flock.

Broilers: Mix 50 g (5 mg / kg) of PROMOTE in a metric ton of feed.

■ **PRESENTATION**

25 kg.

■ **REGISTRATION HOLDER**

VIRBAC RSA (Pty) Ltd
 (Reg No. 1990/003743/07)
 Private Bag X115
 Halfway House
 1685
 South Africa
 Tel: +27 12 657 6000

Namibia Reg. No. V95/25.1/89 NS0