



Prodose® Orange

FOR ANIMAL USE ONLY

Prodose® Orange

Reg. No. G2101 (Act 36/1947)

②+④*

Oral Suspension

Nasalworm, liverfluke, roundworm and milk tapeworm remedy for sheep and goats with residual protection against re-infestation of wireworm and hookworm

CAUTION

STORAGE INSTRUCTIONS

Store at room temperature below 25 °C.

COMPOSITION

Albendazole 1,90 % m/v

Closantel (as sodium) 3,94 % m/v

WARNINGS:

- Do NOT use in milk goats whose milk is intended for human consumption.
- Allow 30 days between last dosing and slaughter for human consumption.
- Handle with care.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS:

- Do not repeat treatments more frequently than every 5 weeks.
- As with all worm remedies it is advisable to ensure animals have been immunised against pulpy kidney disease before drenching.
- Prevent contamination of food, food utensils, etc.
- Wash hands thoroughly after use.
- Do not mix with other products.

DIRECTIONS FOR USE:

Use only as directed

Residues of the compound are blood bound. Meat and organs are virtually free from residues and are therefore suitable for human consumption 30 days after treatment.

SHAKE WELL BEFORE USE.

For oral use only.

Dosage table: 2 ml per 10 kg body mass, e.g.:

Body mass (kg)	Dose (ml)
10	2,0
20	4,0
30	6,0
40	8,0
50	10,0
60	12,0

EFFICACY:

1. **Nasal Worm:** Controls all stages (1st, 2nd and 3rd instar larvae):

2. **Roundworms:**

Prodose Orange controls the following:

	Immature	Adult	Persistency **
Wireworm	*	*	5 weeks
Brown Stomachworm	*	*	-
Bankruptworm	*	*	-
Long-necked Bankruptworm	*	*	-
Hookworm	*	*	2 weeks
Nodular Worm	*	*	-
Large-mouthed Bowelworm	*	*	-

Definition: * Control (≤ 90 % effective), ** Aids in control of (60 - 89 % effective)

3. **Liver Fluke:**

Prodose Orange aids in control of the following:

	6 weeks	Adult
Liver Fluke	**	*
	8 weeks	Adult
Giant Liver Fluke	**	*

Definition: * Control (≤ 90 % effective), ** Aids in control of (60-89 % effective)

4. **Tapeworms: Milk tapeworm (*Moniezia* spp.)**

Class 1 (100 % effective in 80 % of the treated animals)

5. **Ovicidal:** Roundworm eggs present in the animal at the time of dosing are prevented from hatching.

* Benzimidazoles + Salicylanilides

PRESENTATION:

200 ml, 250 ml, 500 ml, 1 kg, 5 kg, 7,5 L, 10 kg, 20 kg and 25 L.

REGISTRATION NUMBER:

VIRBAC RSA (Pty) Ltd
(Reg No. 1990/003743/07)
Private Bag X115
Halfway House
1685, South Africa
Tel: +27 12 657 6000

Namibia Reg. No. V95/18.1.1/42[NS0]

Zimbabwe Reg. No: 2017/80.12.10/9771 (V.M.G.D) 80.10.12 Anthelmintics used in livestock