



# Prodose® Red

FOR ANIMAL USE ONLY

## Prodose® Red

Reg. No. G2102 (Act 36/1947)



Oral Suspension

Roundworm remedy for sheep, goats, cattle and ostriches.

### CAUTION

### STORAGE INSTRUCTIONS

Store at room temperature below 25 °C.

### COMPOSITION

Levamisolehydrochloride 2,56 % m/v

### WARNINGS

- Do not slaughter animals for human consumption within 7 days of treatment.
- Prodose Red should not be dosed to sheep and goats that have not been immunised against pulpy kidney (enterotoxaemia).
- Although Prodose Red can be administered to pregnant animals, care must be exercised in the handling of animals in the late stage of pregnancy.
- In cattle, the odd animal when treated with levamisole may develop an allergic-type reaction of a transient nature which normally disappears within 24 hours. If this is suspected, seek veterinary advice and notify the registration holder.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

### DIRECTIONS FOR USE

#### USE ONLY AS DIRECTED

Use only as a drench. Dose animals according to the following schedule:

**Cattle:** The dose is 15 ml per 50 kg body mass, e.g.:

Body mass (kg)	Dose (ml)
100	30
150	45
200	60
250	75
300	90

**Sheep and goats:** The dose is 3 ml per 10 kg body mass, e.g.:

Body mass (kg)	Dose (ml)
10	3
20	6
30	9
40	12

**Ostriches:** The dose is 6 ml per 5 kg body mass, e.g.:

Body mass (kg)	Dose (ml)
5	6
50	60
150	180

### SAFETY

**Cattle, sheep and goats:** Pregnant ewes and cows, lambs, kids and calves may be dosed with Prodose Red.

### Ostriches:

**Important:** Hold the bird's head up until the dose has been swallowed. Ensure that the dose is given directly into the gullet and not into the windpipe.

### EFFICACY

#### Cattle:

Roundworms	Immatures	Adult
Wireworm	*	*
Cattle Bankruptworm	*	*
Hookworm	-	*
Nodular Worm	**	*
Brown Stomachworm	-	*

Definition: \* Controls (≥ 90 % effective) \*\* Aids in control of (60 – 89 % effective)

#### Sheep and Goats:

Roundworms	Immatures	Adult
Wireworm	*	*
Brown Stomachworm	-	*
Bankruptworm	*	*
Long-necked Bankruptworm	*	*
Hookworm	*	*
Nodular Worm	*	*
Large-mouthed Bowelworm	*	*
Lungworm	**	**

Definition: \* Controls (≥ 90 % effective) \*\* Aids in control of (60 – 89 % effective)

#### Ostriches:

Roundworms	Immatures	Adult
Libyostrongylus douglassii (Wireworm)	**	*
Codiostomum spp.	**	*

Definition: \* Controls (≥ 90 % effective) \*\* Aids in control of (60 – 89 % effective)

<sup>†</sup> Imidazoles

### PRESENTATION:

200 ml, 500 ml, 1 kg, 2 kg, 5 L, 10 kg and 20 kg poly-propylene bottles.

### REGISTRATION NUMBER:

VIRBAC RSA (Pty) Ltd  
(Reg No. 1990/003743/07)  
Private Bag X115  
Halfway House  
1685, South Africa  
Tel: +27 12 657 6000

Namibia Reg. No: V02/18.1.4/8