

## ENDOPARASITICIDES FOR ANIMAL USE ONLY

③<sup>†</sup>

### Prodose<sup>®</sup> Red Soluble Powder

Reg. No. G2166 (Act 36/1947)

#### ROUNDWORM REMEDY FOR SHEEP, GOATS, CATTLE AND OSTRICHES

#### CAUTION

##### ■ STORAGE INSTRUCTIONS

Store at room temperature below 25 °C.

##### ■ COMPOSITION

Levamisole hydrochloride ..... 99 % m/m

##### ■ WARNINGS

- Do not slaughter animals for human consumption within 7 days of treatment.
- No worm remedy, not even PRODOSE RED SOLUBLE POWDER, should be dosed to sheep and goats that have not been immunised against pulpy kidney (enterotoxaemia).
- Although PRODOSE RED SOLUBLE POWDER can be administered to pregnant animals, care must be exercised in the handling of animals in the late stage of pregnancy.
- The odd animal, when treated with levamisole, may develop an allergic type reaction of a transient nature which normally disappears within 24 hours. Should this be severe, consult your veterinarian.  
**Note to veterinarian:** treat symptomatically with an antihistamine. **Under no circumstances must cortisone preparations be used.**
- Keep out of reach of children, uninformed persons and animals.

- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

##### ■ DIRECTIONS FOR USE - Use only as directed

##### Cattle, Sheep and Goats

Dissolve the contents of one sachet (128g) PRODOSE RED SOLUBLE POWDER in 5 litres clean water to obtain a 2,56 % oral solution.

It is advisable to only dissolve sufficient for immediate requirements. Use within 3 days. Remaining solution, after drenching, must not be reused if the solution is cloudy or contains a sediment.

##### Use only as a drench

In cattle use 15 ml per 50 kg body mass. In sheep and goats use 3 ml per 10 kg body mass.

##### Dosage Table

##### Cattle:

Body mass (kg)	Dose (ml)
100	30
150	45
200	65
250	75
300	90

##### Sheep and goats:

Body mass (kg)	Dose (ml)
10	3
20	6
30	9
40	12
50	15

Pregnant ewes and cows, lambs, kids and calves may be dosed with PRODOSE RED SOLUBLE POWDER.

##### Ostriches

Dissolve the contents of one sachet (128 g) PRODOSE RED SOLUBLE POWDER in 2 litres clean water to obtain a 6,4 % oral solution.

It is advisable to only dissolve sufficient for immediate requirements. Use within 3 days. Remaining solution, after drenching, must not be reused if the solution is cloudy or contains a sediment.

##### Use only as a drench

In ostriches use 2,5 ml per 5 kg body mass.

##### Dosage Table

Body mass (kg)	Dose (ml)
5	2,5
50	25
150	75

**Important:** Hold the bird's head up until the dose has been swallowed. Ensure that the dose is given directly into the gullet and not into the windpipe.

##### ■ EFFICACY

##### Internal parasites

For the treatment of gastro-intestinal roundworms in cattle, sheep and goats.

##### Cattle: Roundworm Species

PRODOSE RED SOLUBLE POWDER controls the following:

	Immature	Adult
Wireworm	*	*
Brown Stomachworm		**
Cattle Bankruptworm	*	*
Nodular Worm	**	*
Hookworm		*

Definition: \* Control (≥ 90% effective)

\*\* Aids in the control of (60 – 89 % effective)

##### Sheep and goats: Roundworm Species

PRODOSE RED SOLUBLE POWDER controls the following:

	Immature	Adult
Wireworm	*	*
Brown Stomachworm		*
Bankruptworm	*	*
Long-necked	**	*
Bankruptworm		*
Large-mouthed	*	*
Bowelworm		*
Nodular Worm	*	*
Lungworm	*	*
Hookworm	*	*

Definition: \* Control (≥ 90% effective)

\*\* Aids in the control of (60 – 89 % effective)

##### Ostriches

Highly effective against adult wireworm (*L. douglassii*) and *Codiostomum* worms, also effective against the immature stages.

##### ■ PRESENTATION

One 128 g sachet per unit carton

<sup>†</sup> Imidazoles

##### ■ REGISTRATION HOLDER

VIRBAC RSA (Pty) Ltd  
(Reg No. 1990/003743/07)  
Private Bag X115  
Halfway House  
1685  
South Africa  
Tel: +27 12 657 6000

Namibia Reg. No. V01/18.1.4/124 NSO